XXXIV. *On the Pressure of Vibrations.*


By Lord Rayleigh, *F.R.S.**

THE importance of the consequences deduced by Boltzmann and W. Wien from the doctrine of the pressure of radiation has naturally drawn increased attention to this subject. That æthereal vibrations must exercise a pressure upon a perfectly conducting, and therefore perfectly reflecting, boundary was Maxwell's deduction from his general equations of the electromagnetic field; and the existence of the pressure of light has lately been confirmed experimentally by Lebedew. It seemed to me that it would be of interest to inquire whether other kinds of vibration exercise a pressure, and if possible to frame a general theory of the action.

We are at once confronted with a difference between the conditions to be dealt with in the case of æthereal vibrations and, for example, the vibrations of air. When a plate of polished silver advances against waves of light, the waves indeed are reflected, but the medium itself must be supposed capable of penetrating the plate; whereas in the corresponding case of aerial vibrations the air as well as the vibrations are compressed by the advancing wall. In other cases, however, a closer parallelism may be established. Thus the transverse vibrations of a stretched string, or wire, may be supposed to be limited by a small ring constrained to remain upon the equilibrium line of the string, but capable of sliding freely upon it. In this arrangement the string passes but the vibrations are compressed, when the ring moves inwards.

We will commence with the very simple problem of a pendulum in which a mass C is suspended by a string. B is a ring constrained to the vertical line AD and capable of moving along it; BC$=l$, and θ denotes the angle between BC and AD at any time t. If B is held at rest, BC is an ordinary pendulum, and it is supposed to be executing small vibrations; so that $\theta=\Theta\cos nt$, where $n^2=g/l$. The tension of the string is approximately W, the weight of the bob; and the force tending to push B upwards is at time t W$(1-\cos\theta)$. Now this expression is closely related to the potential energy of the pendulum, for which

Fig. 1.



$$V=Wl(1-\cos\theta).$$

* Communicated by the Author.

The mean upward force upon B is accordingly equal to the mean value of $V \div l$; or since the mean value of V is half the constant total energy E of the system, we conclude that the mean force (L), driving B upwards, is measured by $\frac{1}{2}E/l$.

From the equation

$$L = \tfrac{1}{2}E/l \quad . \quad . \quad . \quad . \quad . \quad . \quad . \quad (1)$$

it is easy to deduce the effect of a *slow* motion upwards of the ring. The work obtained at B must be at the expense of the energy of the system, so that

$$dE = -L\,dl = -\tfrac{1}{2}E\,dl/l.$$

By integration

$$E = E_1 l^{-\frac{1}{2}}, \quad . \quad . \quad . \quad . \quad . \quad . \quad . \quad (2)$$

where E_1 denotes the energy corresponding to $l=1$. From (2) we see that by withdrawing the ring B until l is infinitely great, the whole of the energy of vibration may be abstracted in the form of work done by B, and this by a uniform motion in which no regard is paid to the momentary phase of the vibration.

The argument is nearly the same for the case of a stretched string vibrating transversely in one plane. The string itself may be supposed to be unlimited, while the vibrations are confined by two rings of which one may be fixed and one movable.

If the origin of x be at one end of a string of length l, the transverse displacement may be expressed by

$$y = \phi_1 \frac{\sin \pi x}{l} + \phi_2 \sin \frac{2\pi x}{l} + \ldots, \quad . \quad . \quad . \quad (3)$$

$$\dot{y} = \dot{\phi}_1 \frac{\sin \pi x}{l} + \dot{\phi}_2 \sin \frac{2\pi x}{l} + \ldots, \quad . \quad . \quad . \quad (4)$$

where $\phi_1, \phi_2, \ldots$ are coefficients depending upon the time. For the kinetic and potential energies we have respectively ('Theory of Sound,' § 128)

$$T = \tfrac{1}{4}\rho l \sum_{s=1}^{s=\infty} \dot{\phi}_s^2, \qquad V = \tfrac{1}{4}Wl \sum_{s=1}^{s=\infty} \frac{s^2\pi^2}{l^2}\phi_s^2, \quad . \quad . \quad (5)$$

in which W represents the constant tension and ρ the longitudinal density of the string. For each kind of ϕ the sums of T and V remain constant during the vibration; and the same is of course true of the totals given in (5).

From (3)

$$\frac{dy}{dx} = \frac{\pi}{l}\left(\phi_1 \cos \frac{\pi x}{l} + 2\phi_2 \cos \frac{2\pi x}{l} + \ldots\right),$$

so that when $x=l$

$$\frac{dy}{dx}=\frac{\pi}{l}(-\phi_1+2\phi_2-3\phi_3+\ldots).$$

Accordingly the force tending to drive out the ring at $x=l$ is at time t

$$\tfrac{1}{2}\mathrm{W}\,.\,\frac{\pi^2}{l^2}(-\phi_1+2\phi_2-3\phi_3+\ldots)^2,$$

or in the mean taken over a long interval,

$$\tfrac{1}{2}\mathrm{W}\,.\,\text{Mean}\,\Sigma\frac{s^2\pi^2}{l^2}\phi_s^2.$$

Comparing with (5), we see that the mean force L has the value $2l\times$ mean V; or since mean V = mean T = $\frac{1}{2}$E, E denoting the constant total energy,

$$\mathrm{L}=\mathrm{E}/l. \quad . \quad . \quad . \quad . \quad . \quad . \quad . \quad (6)$$

The force driving out the ring is thus numerically equal to the *longitudinal density of the energy*.

This result may readily be extended to cases where the vibrations are not limited to one plane; and indeed the case in which the plane of the string uniformly revolves is especially simple in that T and V are then constant with respect to time.

If the ring is allowed to move out slowly, we have

$$d\mathrm{E}=-\mathrm{L}\,dl=-\mathrm{E}\,dl/l,$$

or on integration

$$\mathrm{E}=\mathrm{E}_1 l^{-1}, \quad . \quad . \quad . \quad . \quad . \quad . \quad . \quad (7)$$

analogous to (5), though different from it in the power of l involved. If l increase without limit, the whole energy of the vibrations may be abstracted in the form of work done on the ring.

We will now pass on to consider the case of air in a cylinder, vibrating in one dimension and supposed to obey Boyle's law according to which $p=a^2\rho$. By the general hydrodynamical equation ('Theory of Sound,' § 253 *a*),

$$\varpi=\int\frac{dp}{\rho}=-\frac{d\phi}{dt}-\tfrac{1}{2}\mathrm{U}^2, \quad . \quad . \quad . \quad . \quad (8)$$

where ϕ denotes the velocity-potential and U the resultant velocity at any point; so that in the present case, if we integrate over a long interval of time,

$$a^2\int\log p\,dt+\tfrac{1}{2}\int\mathrm{U}^2\,dt \quad . \quad . \quad . \quad . \quad . \quad (9)$$

retains a constant value over the length of the cylinder. If p_0 denote the pressure that would prevail throughout, had there been no vibrations, $p-p_0$ is small and we may replace (9) by

$$a^2\int\left\{\frac{p-p_0}{p_0}-\tfrac{1}{2}\frac{(p-p_0)^2}{p_0^2}\right\}dt+\tfrac{1}{2}\int \mathrm{U}^2\,dt. \quad . \quad . \quad (10)$$

The expression (10) has accordingly the same value at the piston as for the mean of the whole column of length l. Now for the mean of the whole column

$$\int (p-p_0)\,dx=0;$$

and thus if p_1 denote the value of p at the piston where $x=l$,

$$a^2\int\left\{\frac{p_1-p_0}{p_0}-\tfrac{1}{2}\frac{(p_1-p_0)^2}{p_0^2}\right\}dt$$

$$=-\frac{a^2}{2l}\iint\frac{(p-p_0)^2}{p_0^2}\,dx\,dt+\frac{1}{2l}\iint \mathrm{U}^2\,dx\,dt. \quad . \quad (11)$$

It is not difficult to prove that the right-hand member of (11) vanishes. Thus, expressing the motion in terms of ϕ, suppose that

$$\phi=\cos\frac{s\pi x}{l}\cos\frac{s\pi at}{l}. \quad . \quad . \quad . \quad . \quad . \quad . \quad (12)$$

Then

$$p-p_0=\rho_0\,d\phi/dt, \qquad \mathrm{U}=d\phi/dx;$$

and since $p_0=a^2\rho_0$, we get

$$\frac{1}{2l}\iint\left\{\left(\frac{d\phi}{dx}\right)^2-\frac{1}{a^2}\left(\frac{d\phi}{dt}\right)^2\right\}dx\,dt,$$

and this vanishes by (12). Accordingly

$$\int (p_1-p_0)\,dt=\int\frac{(p_1-p_0)^2}{2p_0}dt. \quad . \quad . \quad . \quad . \quad (13)$$

Again by (12)

$$\int\left(\frac{d\phi}{dt}\right)_l^2 dt=\frac{2}{l}\iint\left(\frac{d\phi}{dt}\right)^2 dx\,dt,$$

so that

$$\int (p_1-p_0)\,dt=\frac{1}{p_0 l}\iint (p_1-p_0)^2\,dx\,dt=\frac{\rho_0}{l}\iint \mathrm{U}^2\,dx\,dt.$$

Now $\rho_0\iint \mathrm{U}^2\,dx\,dt$ represents twice the mean total kinetic energy of the vibrations or, what is the same, the constant total energy E. Thus if L denote the mean additional force

due to the vibrations and tending to push the piston out,

$$L = E\, l^{-1}. \quad . \quad . \quad . \quad . \quad . \quad . \quad . \quad . \quad (14)$$

As in the case of the string, the total force is measured by the longitudinal density of the total energy; or, if we prefer so to express it, the additional *pressure* is measured by the volume-density of the energy.

In the last problem, as well as in that of the string, the vibrations are in one dimension. In the case of air there is no difficulty in the extension to two or three dimensions. Thus, if aerial vibrations be distributed equally in all directions, the pressure due to them coincides with *one-third* of the volume-density of the energy. In the case of the string, where the vibrations are transverse, we cannot find an analogue in three dimensions; but a membrane with a flexible and extensible boundary capable of slipping along the surface, provides for two dimensions. If the vibrations be equally distributed in the plane, the force outwards per unit length of contour will be measured by one-half of the superficial density of the total energy.

A more general treatment of the question may be effected by means of Lagrange's theory. If l be one of the coordinates fixing the configuration of a system, the corresponding equation is

$$\frac{d}{dt}\left(\frac{d\mathrm{T}}{dl'}\right) - \frac{d\mathrm{T}}{dl} + \frac{d\mathrm{V}}{dl} = \mathrm{L}, \quad . \quad . \quad . \quad . \quad (15)$$

where T and V denote as usual the expressions for the kinetic and potential energies. On integration over a time t_1

$$\int \frac{\mathrm{L}\,dt}{t_1} = \frac{1}{t_1}\left[\frac{d\mathrm{T}}{dl'}\right] + \frac{1}{t_1}\int\left(\frac{d\mathrm{V}}{dl} - \frac{d\mathrm{T}}{dl}\right)dt.$$

If $d\mathrm{T}/dl'$ remain finite throughout, and if the range of integration be sufficiently extended, the integrated term disappears, and we get

$$\int \frac{\mathrm{L}\,dt}{t_1} = \frac{1}{t_1}\int\left(\frac{d\mathrm{V}}{dt} - \frac{d\mathrm{T}}{dl}\right)dt. \quad . \quad . \quad . \quad (16)$$

On the right hand of (16) the differentiations are partial, the coordinates other than l and all the velocities being supposed constant.

We will apply our equation (16) in the first place to the simple pendulum of fig. 1, l denoting the length of the vibrating portion of the string BC. If x, y be the horizontal

and vertical coordinates of C,

$$x = l \sin\theta, \qquad y = l - l\cos\theta;$$

and accordingly if the mass of C be taken to be unity,

$$\mathrm{T} = \tfrac{1}{2}l'^2(2 - 2\cos\theta) + l'\theta' \,.\, l\sin\theta + \tfrac{1}{2}\theta'^2 l^2, \quad . \quad . \quad (17)$$

l', θ' denoting dl/dt, $d\theta/dt$. Also

$$\mathrm{V} = gl\,(1 - \cos\theta). \quad . \quad . \quad . \quad . \quad . \quad (18)$$

From (17), (18)

$$\frac{d\mathrm{V}}{dl} = g(1 - \cos\theta), \quad \frac{d\mathrm{T}}{dl} = l'\theta' \sin\theta + \theta'^2 l. \quad . \quad . \quad (19)$$

These expressions are general; but for our present purpose it will suffice if we suppose that l' is zero, that is that the ring is held at rest. Accordingly

$$\frac{d\mathrm{V}}{dl} = \frac{\mathrm{V}}{l}, \qquad \frac{d\mathrm{T}}{dl} = \frac{2\mathrm{T}}{l},$$

and (16) gives

$$\int \frac{\mathrm{L}\,dt}{t_1} = \frac{1}{t_1}\int \frac{\mathrm{V} - 2\mathrm{T}}{l}\,dt. \quad . \quad . \quad . \quad . \quad (20)$$

On the right hand of (20) we find the mean values of V and and of T. But these mean values are equal. In fact

$$\int \mathrm{V}\,dt = \int \mathrm{T}\,dt = \tfrac{1}{2}\mathrm{E}t_1, \quad . \quad . \quad . \quad . \quad . \quad (21)$$

if E denote the total energy. Hence, if L now denote the mean value,

$$\mathrm{L} = -\tfrac{1}{2}\mathrm{E}/l, \quad . \quad . \quad . \quad . \quad . \quad . \quad (22)$$

the negative sign denoting that the mean force necessary to hold the ring at rest must be applied in the direction which tends to diminish l, i. e. downwards. In former equations (1), (6), (14), L had the reverse sign.

We will now consider more generally the case of one dimension, using a method that will apply equally whether for example the vibrating body be a stretched string, or a rod vibrating flexurally. All that we postulate is homogeneity of constitution, so that what can be said about any part of the length can be said equally about any other part. In applying Lagrange's method the coordinates are l the length of the vibrating portion, and ϕ_1, ϕ_2, &c. defining, as in (3), the displacement from equilibrium during the vibrations. As functions of l, we suppose that

$$\mathrm{V} \propto l^m, \qquad \mathrm{T} \propto l^n. \quad . \quad . \quad . \quad . \quad . \quad (23)$$

Thus, if L be the force corresponding to l, we get by (16)

$$\int \frac{L\,dt}{t_1} = \frac{1}{t_1}\int\left(\frac{mV}{l} - \frac{nT}{l}\right)dt,$$

in which

$$\int V\,dt = \int T\,dt = \tfrac{1}{2}E\,.\,t_1,$$

E representing as before the constant total energy. Accordingly, L now representing the mean value,

$$L = \frac{(m-n)E}{2l}. \quad . \quad . \quad . \quad . \quad . \quad . \quad . \quad (24)$$

In the case of a medium, like a stretched string, propagating waves of all lengths with the same velocity, $m=-1$, $n=1$, and $L=-E/l$, as was found before.

In the application to a rod vibrating flexurally, $m=-3$, $n=1$, so that

$$L=-2E/l. \quad . \quad . \quad . \quad . \quad . \quad . \quad . \quad (25)$$

If $m=n$, L vanishes. This occurs in the case of the line of disconnected pendulums considered by Reynolds in illustration of the theory of the group velocity *, and the circumstance suggests that L represents the tendency of a group of waves to spread. This conjecture is easily verified. If in conformity with (13) we suppose that

$$V = V_0\, l^m \phi_1^2, \qquad T = T_0\, l^n \dot{\phi}_1^2,$$

and also that

$$\phi_1 = \sin\frac{2\pi t}{\tau}, \qquad \dot{\phi}_1 = \frac{2\pi}{\tau}\cos\frac{2\pi t}{\tau},$$

τ being the period of the vibration represented by the coordinate ϕ_1, we obtain, remembering that the sum of T and V must remain constant,

$$V_0\, l^m = T_0\, l^n\,.\,4\pi/\tau^2.$$

This gives the relation between τ and l. Now v, the wave-velocity, is proportional to l/τ; so that

$$v \propto l^{1-\frac{1}{2}n+\frac{1}{2}m}. \quad . \quad . \quad . \quad . \quad . \quad . \quad . \quad (26)$$

Thus, if u denote the group-velocity, we have by the general theory

$$u/v = \tfrac{1}{2}n - \tfrac{1}{2}m\,; \quad . \quad . \quad . \quad . \quad . \quad . \quad . \quad (27)$$

and in terms of u and v by (24)

$$L = -\frac{uE}{vl}. \quad . \quad . \quad . \quad . \quad . \quad . \quad . \quad (28)$$

* See Proc. Math. Soc. ix. p. 21 (1877); Scientific Papers, i. p. 322. Also Theory of Sound, vol. i. Appendix.

Boltzmann's theory is founded upon the application of Carnot's cycle to the radiation inclosed within movable reflecting walls. If the pressure (p) of a body be regarded as a function of the volume v *, and the absolute temperature θ, the general equation deduced from the second law of thermodynamics is

$$\frac{dp}{d\log\theta} = \mathrm{M}, \quad . \quad . \quad . \quad . \quad . \quad . \quad (29)$$

where $\mathrm{M}\,dv$ represents the heat that must be communicated while the volume alters by dv and $d\theta = 0$. In the application of (29) to radiation we have evidently

$$\mathrm{M} = \mathrm{U} + p, \quad . \quad . \quad . \quad . \quad . \quad . \quad (30)$$

where U denotes the density of the energy—a function of θ only. Hence †

$$\frac{dp}{d\log\theta} = \mathrm{U} + p. \quad . \quad . \quad . \quad . \quad . \quad (31)$$

If further, as for radiation and for aerial vibrations,

$$p = \tfrac{1}{3}\mathrm{U}, \quad . \quad . \quad . \quad . \quad . \quad . \quad . \quad (32)$$

it follows at once that

$$d\log\mathrm{U} = 4\,d\log\theta.$$

whence

$$\mathrm{U} \propto \theta^4, \quad . \quad . \quad . \quad . \quad . \quad . \quad . \quad (33)$$

the well-known law of Stefan. It may be observed that the existence of a pressure is demanded by (31), independently of (32).

If we generalize (32) by taking

$$p = \frac{1}{n}\mathrm{U}, \quad . \quad . \quad . \quad . \quad . \quad . \quad . \quad (34)$$

where n is some numerical quantity, we obtain as the generalization of (33)

$$\mathrm{U} \propto \theta^{n+1}. \quad . \quad . \quad . \quad . \quad . \quad . \quad . \quad (35)$$

It is an interesting question whether any analogue of the second law of thermodynamics can be found in the general theory of the pressure of vibrations, whether for example the energy of the vibrations of a stretched string is partially unavailable in the absence of appliances for distinguishing *phases*. It might appear at first sight that the conclusion already given, as to the possibility of recovering the whole energy by mere retreat of the inclosing ring, was a proof to

* Now with an altered meaning.

† Compare Lorentz, Amsterdam Proceedings, Ap. 1901.

the contrary. This argument, however, will not appear conclusive, if we remember that a like proposition is true for the energy of a gas confined adiabatically under a piston. The residual energy of the molecules may be made as small as we please, but the completion of the cycle by pushing the piston back will restore the molecular energy unless we can first abolish the infinitesimal residue remaining after expansion, and this can only be done with the aid of a body at the absolute zero of temperature. It would appear that we may find an analogue for temperature, so far as the vibrations of *one* system are concerned; but, so far as I can see, the analogy breaks down when we attempt a general theory.

XXXV. *On the Physical Peculiarities of Solutions of Gases in Liquids. By* J. Alfred Wanklyn, *Corresponding Member of the Royal Bavarian Academy of Sciences**.

THOSE who are acquainted with Bunsen's methods of manipulating gases, and especially those chemists who have enjoyed the privilege of Bunsen's personal instruction, will be familiar with Bunsen's zealous care in so using his "Absorptiometer" that every drop of the liquid solvent should come into repeated contact with the gas to be dissolved.

Bunsen's arrangements presuppose that only those particles of the solvent brought into actual contact with the gas become charged with the gas, and that actual passage of the gas from one stratum of liquid to an adjacent stratum either does not take place at all, or is exceedingly slow.

The question may be fairly asked:—Given a quantity of gas in contact with the surface of pure water in a state of absolute quiescence, will that gas penetrate below the surface of the water except with a degree of slowness calling to mind the slow passage of the less diffusive salts which do not traverse a space of 100 millimetres in a fortnight? An experiment on the action of carbonic acid has been made in my laboratory, which I will now describe.

First of all there was the very simple observation that carbonic acid, confined in a tube over mercury, is capable of being absorbed by distilled water kept at rest, that is to say, without being subjected to the shaking up which is usually resorted to in order to bring about such absorptions. The rate of absorption was also noted, and found to be about one

* Communicated by the Author.

THE

LONDON, EDINBURGH, AND DUBLIN

PHILOSOPHICAL MAGAZINE

AND

JOURNAL OF SCIENCE.

CONDUCTED BY

LORD KELVIN, G.C.V.O. D.C.L. LL.D. F.R.S. &c.

JOHN JOLY, M.A. D.Sc. F.R.S. F.G.S.

AND

WILLIAM FRANCIS, PH.D. F.L.S. F.R.A.S. F.C.S.

"Nec aranearum sane textus ideo melior quia ex se fila gignunt, nec noster vilior quia ex alienis libamus ut apes." JUST. LIPS. *Polit.* lib. i. cap. 1. Not.

VOL. III.—SIXTH SERIES.
JANUARY—JUNE 1902.

LONDON:

TAYLOR AND FRANCIS, RED LION COURT, FLEET STREET.

SOLD BY SIMPKIN, MARSHALL, HAMILTON, KENT, AND CO., LD.—T. AND T. CLARK, EDINBURGH;—SMITH AND SON, GLASGOW;—HODGES, FIGGIS, AND CO., DUBLIN;—PUTNAM, NEW YORK;—VEUVE J. BOYVEAU, PARIS;—AND ASHER AND CO., BERLIN.


73149

Q
1
.75

"Meditationis est perscrutari occulta; contemplationis est admirari perspicua .... Admiratio generat quæstionem, quæstio investigationem, investigatio inventionem."—*Hugo de S. Victore.*

——"Cur spirent venti, cur terra dehiscat,
Cur mare turgescat, pelago cur tantus amaror,
Cur caput obscura Phœbus ferrugine condat,
Quid toties diros cogat flagrare cometas,
Quid pariat nubes, veniant cur fulmina cœlo,
Quo micet igne Iris, superos quis conciat orbes
Tam vario motu."

J. B. Pinelli ad Mazonium.



CONTENTS OF VOL. III.

(SIXTH SERIES).

NUMBER XIII.—JANUARY 1902.

	Page
Lord Kelvin on the Clustering of Gravitational Matter in any part of the Universe	1
Dr. W. M. Hicks on the Michelson-Morley Experiment relating to the Drift of the Æther. (Plate I.)	9
Mr. R. F. D'Arcy on the Decomposition of Hydrogen Peroxide by Light, and the Electrical Discharging Action of this Decomposition	42
Mr. Will G. Hormell on the Dielectric Constant of Paraffins.	52
Messrs. R. K. McClung and D. McIntosh on the Absorption of Röntgen Rays by Aqueous Solutions	68
Prof. C. Barus on the Behaviour of the Phosphorus Emanation in Spherical Condensers	80
Mr. H. S. Allen on the Effect of Errors in Ruling on the Appearance of a Diffraction-Grating	92
Lord Kelvin on a New Specifying Method for Stress and Strain in an Elastic Solid	95
Lord Rayleigh: Some General Theorems concerning Forced Vibrations and Resonance	97
Dr. V. H. Veley and Mr. J. J. Manley on the Ionic and Thermal Coefficients of Nitric Acid	118
Mr. B. Hopkinson on the Necessity for Postulating an Æther	123
Mr. H. S. Allen on the Relation between Primary and Secondary Röntgen Radiation	126
Prof. R. W. Wood on the Anomalous Dispersion of Sodium Vapour. (Plates II.–IV.)	128
Mr. Harold Hilton on Capillary Constants of Crystal Faces	144
Prof. J. P. Kuenen and Mr. W. G. Robson on the Vapour-Pressures of Carbon Dioxide and of Ethane at Temperatures below 0° C.	149
Dr. R. A. Lehfeldt on a Voltameter for Small Currents	158

Page

Notices respecting New Books :—
Wilhelm Zenker's Lehrbuch der Photochromie 160
Jacques Hadamard's La Série de Taylor et son prolongement analytique 160

NUMBER XIV.—FEBRUARY.

Mr. W. Sutherland on Ionization, Ionic Velocities, and Atomic Sizes 161
Mr. E. P. Harrison on the Variation with Temperature of the Thermoelectromotive Force, and of the Electric Resistance of Nickel, Iron, and Copper, between the Temperatures of —200° and +1050° 177
Dr. J. C. McLennan on a kind of Radioactivity imparted to certain Salts by Cathode Rays 195
Mr. Harold Hilton: Comparison of various Notations employed in "Theories of Crystal-structure," and a Revision of the 230 Groups of Movements 203
Mr. P. J. Kirkby on the Electrical Conductivities produced in Air by the Motion of Negative Ions 212
Mr. S. H. Burbury on Irreversible Processes and Planck's Theory in relation thereto 225
Dr. J. Buchanan: Note on a Paper by Prof. Fleming, F.R.S., and Mr. Ashton, entitled "On a Model which Imitates the Behaviour of Dielectrics" 240
Dr. P. Chappuis on Gas-Thermometry 243
Mr. G. W. Walker on Asymmetry of the Zeeman Effect .. 247
Notices respecting New Books :—
Expédition Antarctique Belge: Aurores Australes, par M. H. Arctowski 252
Expédition Norvégienne: Resultats des recherches magnétiques, par Kr. Birkeland 252
Prof. H. Weber's Die Partiellen Differential-Gleichungen der Mathematischen Physik 256
The Michelson-Morley Experiment, by Dr. W. M. Hicks .. 256

NUMBER XV.—MARCH.

Page
Lord Kelvin: Aepinus Atomized 257
Mr. R. K. McClung on the Rate of Recombination of Ions in Gases under Different Pressures..... 283
Dr. F. G. Donnan on the Condensation of the Vapours of Organic Liquids in Presence of Dust-free Air. (Plate V.) 305
Prof. T. R. Lyle on Circular Filaments or Circular Magnetic Shells equivalent to Circular Coils, and on the equivalent Radius of a Coil. (Plate VI.)..... 310
Prof. A. A. Michelson on the Velocity of Light 330
Lord Rayleigh on the Pressure of Vibrations..... 338
Mr. J. A. Wanklyn on the Physical Peculiarities of Solutions of Gases in Liquids 346
Mr. Clive Cuthbertson on the Arrangement of Bands in the First Group of the Positive Band-Spectrum of Nitrogen .. 348
Prof. J. J. Thomson on the Effect of a Transverse Magnetic Field on Metallic Resistance 353
Notices respecting New Books :—
Original Papers by the late John Hopkinson, D.Sc., F.R.S. 357
Dr. J. A. Fleming's Handbook for the Electrical Laboratory and Testing-Room 358
Annuaire du Bureau des Longitudes 359
On the Fluorescence and Absorption-Spectrum of Sodium Vapour, by Prof. R. W. Wood 359

NUMBER XVI.—APRIL.

Mr. N. E. Gilbert: Some Experiments upon the Relations between Æther, Matter, and Electricity..... 361
Mr. E. B. H. Wade on a new Hygrometric Method..... 380
Dr. E. H. Barton and Mr. S. C. Laws on Air-Pressures used in playing Brass Instruments. (Plates VII. & VIII.) .. 385
Prof. J. Trowbridge on the Induction Coil 393
Prof. R. W. Wood on a suspected case of the Electrical Resonance of Minute Metal Particles for Light-waves. A New Type of Absorption 396
Dr. T. Muir on the Aggregates of Minors of an Axisymmetric Determinant 410
Lord Rayleigh on the Question of Hydrogen in the Atmosphere 416
Dr. J. W. Mellor on a Law of Molecular Attraction 423
Mr. C. A. Chant on an Experimental Investigation into the 'Skin'-effect in Electrical Oscillators..... 425

Page
Lord Kelvin on a New Specifying Method for Stress and Strain in an Elastic Solid 444
Prof. W. Cassie on Multiple Transmission Fixed-arm Spectroscopes 449
Notices respecting New Books :—
Otto Koll's Die Theorie der Beobachtungsfehler und die Methode der kleinsten Quadrate mit ihrer Anwendung auf die Geodäsie und die Wassermessungen 458
Proceedings of the Geological Society :—
Mr. T. Codrington on a Submerged and Glaciated Rock-Valley recently exposed to view in Carmarthenshire.. 458
Mr. F. Rutley on an Altered Siliceous Sinter from Builth (Brecknockshire) 459
Dr. Nils Ekholm on the Meteorological Conditions of the Pleistocene Epoch 459
Mr. H. B. Stocks on the Origin of certain Concretions in the Lower Coal-Measures 460

NUMBER XVII.—MAY.

Mr. J. E. Petavel on a Recording Manometer for High-Pressure Explosions. (Plates IX. & X.) 461
Prof. S. R. Cook on Flutings in a Sound-Wave and the Forces due to a Flux of a Viscous Fluid around Spheres 471
Prof. J. D. Everett on Focal Lines, and Anchor-Ring Wave-Fronts 483
Prof. T. R. Lyle and Mr. R. Hosking on the Temperature Variations of the Specific Molecular Conductivity and of the Fluidity of Sodium Chloride Solutions. (Plates XI. & XII.) 487
Mr. J. A. Wanklyn on the Physical Peculiarities of Solutions of Gases in Liquids.—No. 2 498
Dr. W. Mansergh Varley on the Magnetism induced in Iron by Rapidly Oscillating Current-fields 500
Prof. P. Lewis on the Rôle of Water Vapour in Gaseous Conduction 512
Mr. W. Williams on the Temperature Variation of the Electrical Resistances of Pure Metals, and Allied Matters . 515
Dr. J. C. Beattie and Mr. J. T. Morrison on the Magnetic Elements at the Cape of Good Hope from 1605 to 1900.. 532
Prof. K. Olszewski on the Determination of Inversion Temperature of Kelvin Effect in Hydrogen 535
Mr. J. Walker on MacCullagh and Stokes's Elliptic Analyser, and other Applications of a Geometrical Representation of the State of Polarization of a Stream of Light 541

Page

Proceedings of the Geological Society :—
Prof. T. T. Groom on the Sequence of the Cambrian and Associated Beds of the Malvern Hills 550
Lieut.-Col. T. English on Coal and Petroleum-Deposits in European Turkey 551
Prof. J. W. W. Spencer on the Geological and Physical Development of Dominica, with Notes on Martinique, St. Lucia, St. Vincent, and the Grenadines 552
Prof. J. W. W. Spencer on the Geological and Physical Development of Barbados, with Notes on Trinidad .. 553
The Michelson-Morley Experiment, by William B. Cartmel; with Note by Dr. W. M. Hicks 555

NUMBER XVIII.—JUNE.

Prof. J. S. Townsend on the Conductivity produced in Gases by the aid of Ultra-Violet Light 557
Mr. F. L. Hitchcock on Vector Differentials..... 576
Prof. J. A. Pollock and Mr. O. U. Vonwiller: Some Experiments on Electric Waves in Short Wire Systems, and on the Specific Inductive Capacity of a Specimen of Glass.. 586
Prof. R. W. Wood on the Absorption, Dispersion, and Surface-Colour of Selenium 607
Prof. J. P. Kuenen and Mr. W. G. Robson on the Thermal Properties of Carbon Dioxide and of Ethane..... 622
Mr. A. E. Tutton on the Thermal Expansion of Porcelain .. 631
Mr. J. Patterson on the Change of the Electrical Resistance of Metals when placed in a Magnetic Field 643
Notices respecting New Books :—
Dr. A. Korn's Lehrbuch der Potentialtheorie. Part II. 657
Dr. H. W. Bakhuis Roozeboom's Die Heterogenen Gleichgewichte vom Standpunkte der Phasenlehre .. 657
Prof. H. C. Jones's Outlines of Electrochemistry 658
Halbmonatliches Litteraturverzeichniss der Fortschritte der Physik im Jahre 1902 658
The Production of a Magnetic Field by a Flight of Charged Particles, by Prof. R. W. Wood..... 659

Index 660

PLATES.

I. Illustrative of Dr. W. M. Hicks's Paper on the Michelson-Morley Experiment relating to the Drift of the Æther.

II.–IV. Illustrative of Prof. R. W. Wood's Paper on the Anomalous Dispersion of Sodium Vapour.

V. Illustrative of Dr. F. G. Donnan's Paper on Condensation of the Vapours of Organic Liquids in Dust-free Air.

VI. Illustrative of Prof. T. R. Lyle's Paper on Circular Filaments or Circular Magnetic Shells equivalent to Circular Coils.

VII. & VIII. Illustrative of Dr. E. H. Barton and Mr. S. C. Laws's Paper on Air-Pressures used in playing Brass Instruments.

IX. & X. Illustrative of Mr. J. E. Petavel's Paper on a Recording Manometer for High-Pressure Explosions.

XI. & XII. Illustrative of Prof. T. R. Lyle and Mr. R. Hosking's Paper on the Temperature Variations of the Specific Molecular Conductivity and of the Fluidity of Sodium Chloride Solutions.